

The current optimal results with these developments are as follows: Using a pressure infusion bag at 300 mm of mercury, standard IV tubing and an 8.5 French catheter, in vivo flow rates of 500 ml per minute are achieved. Experimentally, using the same catheter with 6.4 mm tubing under 600 mm of mercury pressure, flow rates in vitro of crystalloid exceeded 3,000 ml per minute!

A word of warning: In the confusion surrounding the resuscitation of an exsanguinating patient, massive volume overload becomes a real danger. Therefore, careful ongoing evaluation and monitoring of intravascular volume are mandatory.

ROBERT H. DAILEY, MD
Oakland, California

REFERENCES

- Dailey RH: Femoral vein cannulation: A review. *J Emerg Med*, in press
Dailey RH: Use of wire-guided (Seldinger-type) catheters in the emergency department. *Ann Emerg Med* 1983 Aug; 12:489-492
Haynes BE, Carr FJ, Niemann JT: Catheter introducers for rapid fluid resuscitation. *Ann Emerg Med* 1983 Oct; 12:606-609
Iserson KV, Reeter AK: Rapid fluid replacement: A new methodology. *Ann Emerg Med* 1984 Feb; 13:97-100
Mateer JR, Thompson BM, Tucker J, et al: Effects of high pressure and large-bore tubing on IV flow rates. *Ann Emerg Med (UAEM Abstracts, 14th Annual Meeting)* 1984 May; 13:405-406

Diagnosing Cases of Ectopic Pregnancy

ECTOPIC PREGNANCY is the leading cause of maternal death in the first trimester of pregnancy. The total number of cases of ectopic pregnancy reported in the United States rose from less than 18,000 in 1970 to greater than 52,000 in 1980. Ectopic pregnancy is extremely difficult to diagnose clinically. More than half of all cases of ectopic pregnancy are missed at first presentation, more than two thirds of ectopic pregnancies are ruptured at the time of diagnosis and 10% to 20% of patients suffer hypovolemic shock before definitive treatment is obtained.

Ectopic pregnancy must be suspected in all women of childbearing age presenting with either abdominal pain, irregular vaginal bleeding or both. While physical examination alone is unreliable in diagnosing ectopic pregnancy, culdocentesis has been found to be positive in from 75% to 94% of cases.

The two-minute agglutination-inhibition slide test for human chorionic gonadotropin (HCG) in urine remains negative until more than 1,500 mIU of HCG per ml is present, and from 30% to 50% of cases of ectopic pregnancy will be missed. The two-hour urine slide test is more sensitive (positive at greater than 250 mIU HCG per ml) but as much as 10% to 20% of cases of ectopic pregnancy may still be

missed. Qualitative serum β -HCG assays are currently available, inexpensive, require only 30 minutes to do and give a positive report at a serum level of greater than 35 mIU HCG per ml. In a recent review of the literature, this test was positive in 439 of 445 (98.7%) cases of ectopic pregnancy. Quantitative serum β -HCG assays measuring levels less than 25 mIU per ml exist but are not widely available and require much longer to do.

Real-time ultrasonography is of great help in establishing the diagnosis of unruptured ectopic pregnancy. Its primary role lies in documenting a normal intrauterine pregnancy at about five to six weeks of gestation. Such a finding essentially excludes the possibility of ectopic pregnancy because the incidence of coexisting ectopic pregnancy and intrauterine pregnancy is about 1 in 30,000 pregnancies. Ultrasound examination may be of secondary importance in supporting a diagnosis of possible ectopic pregnancy by showing an adnexal mass or fluid within the cul-de-sac, or both. The ability to identify an adnexal mass as an ectopic pregnancy rather than a large ovarian cyst, hydrosalpinx, tubo-ovarian abscess or other cause of adnexal enlargement varies from center to center.

When a case of ectopic pregnancy is suspected, culdocentesis should be done. Positive findings on culdocentesis warrant a gynecologic consultation. A negative or nondiagnostic culdocentesis should be followed by pregnancy testing. A rapid two-minute urine pregnancy test that is positive should be followed by pelvic ultrasound examination. An ultrasound examination that does not show a normal intrauterine gestational sac should be followed by laparoscopy. If the two-minute urine pregnancy test is negative, it should be followed by a serum β -HCG test. A negative β -HCG test rules out an ectopic pregnancy in all but about 1% of cases. A positive β -HCG test should be followed by pelvic ultrasound study or laparoscopy. In one recently published study in which such an approach was used, all 16 patients with ectopic pregnancy from a study group of 81 patients suspected of having the disease were successfully identified.

ROBERT S. HOCKBERGER, MD
Torrance, California

REFERENCES

- Bryson SC: Beta-subunit of human chorionic gonadotropin, ultrasound, and ectopic pregnancy: A prospective study. *Am J Obstet Gynecol* 1983 May 15; 146:163-165
Gleicher N, Giglia RV, Deppe G, et al: Direct diagnosis of unruptured ectopic pregnancy by real-time ultrasonography. *Obstet Gynecol* 1983 Apr; 61:425-428
Olson CM, Holt JA, Alenghat E, et al: Limitations of qualitative serum beta-HCG assays in the diagnosis of ectopic pregnancy. *J Reprod Med* 1983 Dec; 28:838-842
Portuondo JA, Remacha MJ, Llaguno MR: Ectopic pregnancy early diagnosis limitations. *Int J Gynaecol Obstet* 1982 Oct; 20:371-378

ADVISORY PANEL TO THE SECTION ON EMERGENCY MEDICINE

CHARLES J. FISHER, JR, MD

Advisory Panel Chair

CMA Scientific Board Representative

University of California, Davis

MICHAEL DORSO, MD
CMA Section Chair
Roseville
DAVID KALIFON, MD
CMA Section Secretary
Malibu
JOHN R. TUCKER, MD
CMA Section Assistant Secretary
Sacramento

DONALD MINESINGER, MD
Loma Linda University
MICHAEL ELIASTAM, MD
Stanford University
KYM SALNESS, MD
University of California, Irvine
JEROME HOFFMAN, MD
Section Editor
University of California, Los Angeles

WILLIAM BAXT, MD
University of California, San Diego
FRANK LEWIS, MD
University of California, San Francisco
GAIL ANDERSON, MD
University of Southern California
DENIS J. KOLLAR, MD
West Covina

MICHAEL BRESLER, MD
San Francisco
LARRY J. BARAFF, MD
Los Angeles
LARRY BEDARD, MD
Sausalito
ELLEN TALIAFERRO, MD
Santa Monica